

We claim:

1. A method for inducing an immunological response against a malignant pancreatic cell in an individual, said method comprising the steps of: selecting an individual having malignant pancreatic cells or at risk for developing such a pancreatic tumor, administering to the individual a first vector containing a first gene, or antigenic portion thereof, that encodes a pancreatic tumor-associated antigen (PTAA), and at regular intervals thereafter administering at least a second vector containing a gene encoding PTAA or antigenic portion thereof, wherein if CEA or MUC-1 or an antigenic portion thereof or a modified version thereof is the PTAA, there must be a second PTAA present.
2. The method of claim 1, further comprising administering granulocyte-macrophage colony stimulating factor (GM-CSF).
3. The method of claim 1 or 2, further comprising administering at least one co-stimulatory molecule.
4. The method of claim 3, wherein the co-stimulatory molecule is administered as a gene contained within the same vector as the vector containing gene encoding the PTAA.
5. The method of claim 3, wherein the co-stimulatory molecule is administered as a gene contained within a different vector than the vector containing gene encoding the PTAA.
6. The method according to claims 1-5, wherein said PTAA, or antigenic portion thereof, is contained in a poxvirus vector.
7. The method according to claim 6, wherein said poxvirus vector is selected from the group consisting of an orthopox virus vector; avipox virus vector; a suipox virus vector; a capripox virus vector; a leporipox virus vector; and an iridovirus vector.
8. The method according to claim 6, wherein said pox virus vector is a replication impaired or non-replicating pox virus vector.
9. The method according to claim 8, wherein said pox virus vector is an orthopox vector.
10. The method according to claim 9, wherein said orthopox virus vector is vaccinia.

11. The method of claims 1-10, wherein the PTAA is selected from the group consisting of carcinoembryonic antigen, mucin, ras, gastrin, erbB2, interferon $\alpha$ , tumor necrosis factor- $\alpha$ , hMP-9 immunotoxin, antigenic portions thereof, and modified versions thereof.
12. The method of claim 11, wherein the PTAA is a mucin selected from the group consisting of MUC-1, MUC-2, MUC-3, MUC-4, MUC-5AC, MUC-5B, MUC-6, MUC-7, MUC-11, MUC-12, and antigenic portions thereof and modified versions thereof.
13. The method of claim 11, wherein the modified version thereof is wobbled-MUC.
14. The method of claims 1-11, wherein the vectors contain genes encoding at least two PTAAAs or antigenic portions thereof.
15. The method of claim 14, wherein the PTAA is CEA and a mucin.
16. The method of claim 15, wherein the mucin is wobbled MUC-1 or wobbled mini-MUC.
17. The method of claims 1-16, wherein one to three administrations at set intervals are made by an orthopox vector containing the at least one PTAA or antigenic portion thereof and multiple administrations at set intervals are made by an avipox vector containing the at least one PTAA or antigenic portion thereof.
18. The method of claim 17, wherein the orthopox vector is vaccinia.
19. The method of claim 18, wherein the vaccinia is an attenuated vaccinia.
20. The method of claim 19, wherein the attenuated vaccinia is MVA or NYVAC.
21. The method of claims 17, 18, 19 or 20, wherein the orthopox vector is administered before the avipox vector is administered.
22. The method of claim 21, wherein the set interval is 20 days to 90 days.
23. A kit for enhancing a protective immune reaction against a pancreatic tumor comprising at least one pox vector encoding at least two PTAAAs or antigenic portion thereof.
24. The kit of claim 23, wherein one PTAA is CEA and the at least second PTAA is MUC-1 or mini-MUC.

25. The kit of claim 24, wherein mini-MUC contains six tandem repeats, is wobbled at the nucleic acid level to prevent excision during homologous recombination, and may further contain additional mutations to enhance immunogenicity.